

# Spacetimes with a separable Klein–Gordon equation in higher dimensions

Ivan Kolář\* and Pavel Krtouš†

*Institute of Theoretical Physics, Faculty of Mathematics and Physics,  
Charles University in Prague, V Holešovičkách 2, 180 00 Prague, Czech Republic*

(Dated: September 29, 2015)

We study spacetimes that lead to a separable Klein–Gordon equation in a general number of dimensions. We introduce an ansatz for the metric in higher dimensions motivated by analogical work by Carter in four dimensions and find solutions of the Klein–Gordon equation. For such a metric we solve the Einstein equations and regain the Kerr–NUT–(A)dS spacetime as one of our results. Other solutions lead to the Einstein–Kähler metric of a Euclidean signature. Next we investigate a warped geometry of two Klein–Gordon separable spaces with a properly chosen warped factor. We show that the resulting metric leads also to a separable Klein–Gordon equation and we find the corresponding solutions. Finally, we solve the Einstein equations for the warped geometry and obtain new solutions.

PACS numbers: 04.50.Gh, 02.30.Ik, 04.70.Bw

## I. INTRODUCTION

The Kerr solution in four dimensions is one of the most important solutions of the Einstein equations. It allows us to describe the exterior of a stationary rotating central source. Although it deals with a rather elementary physical configuration, it took a significant effort and rather long time to find it—until 1963, 47 years after the discovery of the spherical Schwarzschild solution [1].

The Kerr solution is not important only for its astrophysical applications but it is also exceptional for its mathematical structure and high symmetry. Geodesic motion in this spacetime is completely integrable and the main physical field equations can be solved by separation of constants. Exactly these properties are the cornerstones of Carter’s rederivation and generalization of the Kerr solution in 1968. In his seminal paper [2], cf. also [3], he studied spaces allowing a separable Hamilton–Jacobi equation and Klein–Gordon equation (referred to as the Schrödinger equation by Carter), which satisfy the Einstein equations. He obtained the Kerr solutions with a cosmological constant and NUT parameter, and some other related spaces. The aim of this paper is to follow this route in a higher-dimensional setting.

Higher dimensional solutions of the Einstein equations have become popular in recent years due to their significance in string theory or in the AdS/CFT context. The study of general relativity in a general number of dimensions is interesting also just from the mathematical point of view, and it can reveal some important properties of various structures and equations. Similarly to the four dimensions, finding a solution describing a generally rotating black hole in higher dimensions was a challenging task. The spherically symmetric Tangherlini–Schwarzschild solution [4] was found in the same year as the Kerr solution, in prehistory of the higher-dimensional

studies. The solution describing a rotating asymptotically flat black hole was discovered by Myers and Perry [5] in 1986. Generalization for a nontrivial cosmological constant has been studied on various levels of generality [6–8] until 2006 when Chen, Lü, and Pope were able to add also the so-called NUT parameters and to write down the most general metric, describing the geometry known today as the *Kerr–NUT–(A)dS spacetime* [9, 10].

It is remarkable that although the derivation of this geometry occurred through an unrelated route (the Kerr–Schild form of the metric) the final form of the metric is a straightforward generalization of Carter’s four-dimensional Kerr–NUT–(A)dS metric [2]. It turns out also that most of the miraculous properties of the Kerr solution survive in higher dimensions: geodesic motion is completely integrable [11], Hamilton–Jacobi, Dirac, and Klein–Gordon equations are separable [12–15], and the geometry possesses a high degree of symmetry encoded in the existence of a tower of Killing vectors and tensors [16, 17]. These results have completed a series of previous works on integrability and separability of the basic physical equations in black hole spacetimes in particular dimensions and for particular values of spacetime parameters [18–33]. It turns out that the Kerr–NUT–(A)dS spacetime is a very prominent example of a fully integrable and separable system [34, 35].

It is interesting that all these properties hold for a wider class of metrics—the so-called off-shell Kerr–NUT–(A)dS spaces—obtained by relaxing conditions on some of the metric functions in the Kerr–NUT–(A)dS solution. It was shown that this class is uniquely determined by requiring the existence of a principal closed conformal Killing–Yano form [36–39].

It is natural to ask if the elegant Carter’s method could be employed also in higher dimensions and whether it could not lead to a more general class of solutions. In this paper we investigate this possibility. We assume a metric form motivated by a combination of Carter’s metric in four dimensions and the Kerr–NUT–(A)dS metric in a general number of dimensions. We show that such a

\* Ivan.Kolar@utf.mff.cuni.cz

† Pavel.Krtous@utf.mff.cuni.cz

metric leads to a separable Klein–Gordon equation and we find the corresponding solutions. We call this class the Klein–Gordon simple separable metrics.<sup>1</sup> Moreover, according to [40], it seems that metrics which yield a separable Hamilton–Jacobi equation should also have this form.

The introduced class generalizes the off-shell Kerr–NUT–(A)dS metrics by introducing one functional reparametrization freedom for each latitude and radial coordinate. It is indeed a wider class, and thus not all Klein–Gordon simple separable metrics possess the (standard) principal closed conformal Killing–Yano form, however, they still allow the tower of Killing tensors and vectors. In fact, the class of simple separable metrics belongs to a wider class of metrics admitting a generalized closed conformal Killing–Yano form with torsion [41], which justifies the exceptional symmetry.

Next, we solve the Einstein equations in the class of Klein–Gordon simple separable metrics. One family of solutions contains exactly the on-shell Kerr–NUT–(A)dS spacetimes. However, we also find another family of solutions, which are special Einstein–Kähler metrics. Unfortunately, these metrics necessarily have a Euclidean signature, which cannot be Wick-rotated to the physical sector. Probably, for this reason, a similar solution was not further considered by Carter in four dimensions.

However, in higher dimensions, we have several ways of building a geometry from smaller blocks. One of such methods is a warped product of two metrics. We employ this method and investigate the warped product of two off-shell Klein–Gordon simple separable metrics. We prove that, although such a metric does not fit in the Klein–Gordon simple separable class, it leads again to a separable Klein–Gordon equation.

For this metric we solve the Einstein equations and prove that the component of the warped product under the warped factor must be the on-shell Klein–Gordon simple separable metric. Since this component must be Euclidean (temporal direction belongs to the other component), it is either an on-shell Kerr–NUT–(A)dS Euclidean instanton (a deformed, twisted sphere, cf. [42]) or the on-shell Klein–Gordon simple separable Einstein–Kähler metric. The former solution has been recently obtained as a limit of Kerr–NUT–(A)dS spacetime where several rotations are sent to zero with properly scaled coordinates, mass, and NUT charges [42]. It also has been studied in the context of warped products of Kerr–NUT–(A)dS spaces [43]. The latter case is, to our knowledge, a new solution of the Einstein equations in higher dimensions.

The paper is organized as follows. In Sec. II we introduce the class of Klein–Gordon simple separable metrics

and discuss some of their properties. In particular, we show that such a metric possesses Killing vectors and rank-two Killing tensors, which enable us to solve the Klein–Gordon equation by separation of variables. Then, we solve the Einstein equations and, thus, obtain the on-shell Klein–Gordon simple separable spacetimes.

Section III is devoted to the warped spacetimes. We briefly summarize the main properties of general warped spaces. Next, we consider a warped geometry of two Klein–Gordon simple separable metrics with a suitable chosen warped factor. Finally, adopting the warped metric ansatz, we find two families of solutions of the Einstein equations.

We conclude with a brief summary in Sec. IV. In Appendixes A and B, we gather some useful identities and expressions for the curvature of the studied metrics.

For simplicity, we restrict our discussion to even dimensions. Generalization to odd dimensions is straightforward, but one has to deal with additional “odd” terms in most expressions.

## II. KLEIN–GORDON SIMPLE SEPARABLE SPACETIMES

### A. Off-shell metric

Consider a metric of even dimension<sup>2</sup>  $D = 2N$ ,

$$g = \sum_{\mu} \left[ \frac{U_{\mu}}{X_{\mu}} (\mathbf{d}x_{\mu})^2 + \frac{X_{\mu}}{U_{\mu}} \left( \sum_j A_{\mu}^{(j)} \mathbf{d}\psi_j \right)^2 \right], \quad (2.1)$$

where  $X_{\mu} = X_{\mu}(x_{\mu})$  are arbitrary functions of a single variable  $x_{\mu}$ . Functions  $U_{\mu}$  and  $A_{\mu}^{(i)}$  are defined by

$$U_{\mu} = \prod_{\substack{\nu \\ \nu \neq \mu}} (Z_{\nu} - Z_{\mu}), \quad A_{\mu}^{(i)} = \sum_{\substack{\nu_1, \dots, \nu_i \\ \nu_1 < \dots < \nu_i \\ \nu_k \neq \mu}} Z_{\nu_1} \dots Z_{\nu_i}. \quad (2.2)$$

We set  $U_1 = 1$  for  $N = 1$  and  $A_{\mu}^{(0)} = 1$ . For further reference, we also define functions

$$U = \prod_{\substack{\mu, \nu \\ \mu < \nu}} (Z_{\mu} - Z_{\nu}), \quad A^{(i)} = \sum_{\substack{\nu_1, \dots, \nu_i \\ \nu_1 < \dots < \nu_i}} Z_{\nu_1} \dots Z_{\nu_i}, \quad (2.3)$$

where we set  $U = 1$  for  $N = 1$  and  $A^{(0)} = 1$ . Here,  $Z_{\mu} = Z_{\mu}(x_{\mu})$  are arbitrary functions of a single variable  $x_{\mu}$ . In what follows, we also assume that all functions  $Z_{\mu}$  are functionally independent, i.e.,  $Z'_{\mu} \neq 0$ . However, some results can be generalized to the cases

<sup>1</sup> The class is specified by the ansatz (2.1) below. Although all such metrics lead to a separable Klein–Gordon equation, not all metrics leading to a separable Klein–Gordon equation belong to this class, as we will observe below.

<sup>2</sup> In what follows, we do not assume an implicit sum over greek indices  $\mu, \nu, \dots$  and latin indices  $i, j, \dots$ . Unless otherwise stated, indices have ranges  $\mu, \nu = 1, \dots, N$  and  $i, j = 0, \dots, N-1$ , respectively. We use here shortened notations  $\sum_{\mu} \equiv \sum_{\mu=1}^N$ ,  $\sum_i \equiv \sum_{i=0}^{N-1}$  and  $\prod_{\mu} \equiv \prod_{\mu=1}^N$ ,  $\prod_i \equiv \prod_{i=0}^{N-1}$ .

when  $Z_\mu = r_\mu$ , where  $r_\mu$  are constants satisfying  $r_\mu \neq r_\nu$ ,  $\mu \neq \nu$ .

Metric (2.1) generalizes the four-dimensional metric introduced by Carter in [2, 3], and since it leads also to the separable Klein–Gordon equation, we call it the Klein–Gordon simple separable metric.

It is also clearly motivated by the (off-shell) Kerr–NUT–(A)dS metric in even dimensions [9, 10], which can be obtained by setting  $Z_\mu = x_\mu^2$ . We could also rewrite the metric in an alternative form,

$$g = \sum_\mu \left[ \frac{U_\mu}{Y_\mu} (\mathbf{d}x_\mu)^2 + \frac{X_\mu}{U_\mu} \left( \sum_j A_\mu^{(j)} \mathbf{d}\psi_j \right)^2 \right], \quad (2.4)$$

in which  $Z_\mu$  is eliminated by introducing a new coordinate  $x_\mu$  given exactly by relation  $Z_\mu = x_\mu^2$ ; the freedom in  $Z_\mu$  is then shifted to a suitable defined new metric function  $Y_\mu = Y_\mu(x_\mu)$ . In this form, the metric is clearly of type A of [41], which admits a generalized principal Killing–Yano tensor with torsion. The metric (2.4) was also used in studying a higher-dimensional generalization of the Wahlquist metric [44].<sup>3</sup> In the following, we will use the original form (2.1) of the metric.

The determinant  $\mathfrak{g}$  of the metric (2.1) in coordinates  $x_\mu, \psi_k$  reads

$$\mathfrak{g} = U^2. \quad (2.5)$$

We introduce the orthonormal frame  $e_\mu, \hat{e}_\mu$ , and dual covector frame  $e^\mu, \hat{e}^\mu$ ,

$$\begin{aligned} e_\mu &= \sqrt{\frac{X_\mu}{U_\mu}} \frac{\partial}{\partial x_\mu}, \quad \hat{e}_\mu = \sqrt{\frac{U_\mu}{X_\mu}} \sum_k \frac{(-Z_\mu)^{N-1-k}}{U_\mu} \frac{\partial}{\partial \psi_k}, \\ e^\mu &= \sqrt{\frac{U_\mu}{X_\mu}} \mathbf{d}x_\mu, \quad \hat{e}^\mu = \sqrt{\frac{X_\mu}{U_\mu}} \sum_k A_\mu^{(k)} \mathbf{d}\psi_k. \end{aligned} \quad (2.6)$$

The Riemann tensor of metric (2.1) is given in (B4) in Appendix. The Ricci tensor can be written in the form

$$\begin{aligned} \text{Ric} &= \sum_\mu T_\mu e^\mu e^\mu + \sum_\mu \left[ T_\mu + \frac{1}{2} \sum_{\substack{\nu \\ \nu \neq \mu}} \frac{S_{\nu,\mu}}{Z'_\mu} \frac{X_\mu}{U_\mu} \right] \hat{e}^\mu \hat{e}^\mu \\ &+ \sum_{\substack{\mu,\nu \\ \nu \neq \mu}} \frac{1}{2} \frac{S_\mu - S_\nu}{Z_\mu - Z_\nu} \sqrt{\frac{X_\mu}{U_\mu} \frac{X_\nu}{U_\nu}} \hat{e}^\mu \hat{e}^\nu, \end{aligned} \quad (2.7)$$

where<sup>4</sup>

$$\begin{aligned} S_\mu &= Z''_\mu + \sum_{\substack{\kappa \\ \kappa \neq \mu}} \frac{Z'_\mu{}^2 - Z'_\kappa{}^2}{Z_\mu - Z_\kappa}, \\ T_\mu &= -\frac{1}{2} \frac{X''_\mu}{U_\mu} - \frac{1}{2} \sum_{\substack{\nu \\ \nu \neq \mu}} \frac{S_{\mu,\nu}}{Z'_\nu} \frac{X_\nu}{U_\nu} \\ &+ \frac{1}{2} \sum_{\substack{\nu \\ \nu \neq \mu}} \frac{1}{U_\mu} \left[ \frac{Z'_\mu X'_\mu - Z''_\mu X_\mu}{Z_\mu - Z_\nu} + \frac{Z'_\nu X'_\nu - Z''_\nu X_\nu}{Z_\nu - Z_\mu} \right]. \end{aligned} \quad (2.8)$$

The scalar curvature reads

$$\mathcal{R} = - \sum_\mu \frac{X''_\mu}{U_\mu}. \quad (2.9)$$

The metric (2.1) admits a generalized closed conformal Killing–Yano 2-form with torsion [41]

$$h = \sum_\mu \sqrt{Z_\mu} e^\mu \wedge \hat{e}^\mu, \quad (2.10)$$

which guarantees the existence of hidden symmetries encoded by rank-two Killing tensors,

$$\begin{aligned} {}^i k &= \sum_\mu A_\mu^{(i)} \left[ \frac{X_\mu}{U_\mu} \left( \frac{\partial}{\partial x_\mu} \right)^2 + \frac{U_\mu}{X_\mu} \left( \sum_k \frac{(-Z_\mu)^{N-1-k}}{U_\mu} \frac{\partial}{\partial \psi_k} \right)^2 \right] \\ &= \sum_\mu A_\mu^{(i)} (e_\mu e_\mu + \hat{e}_\mu \hat{e}_\mu). \end{aligned} \quad (2.11)$$

The geometry (2.1) also possesses a set of explicit symmetries given by the Killing vectors,

$${}^i l = \frac{\partial}{\partial \psi_i} = \sum_\mu \sqrt{\frac{X_\mu}{U_\mu}} A_\mu^{(i)} \hat{e}_\mu. \quad (2.12)$$

Since the operators (2.13) constructed in the next subsection mutually commute, the corresponding phase-space observables  ${}^j k^{ab} p_a p_b$ ,  ${}^k l^a p_a$  (quadratic and linear in momentum  $\mathbf{p}$ ) must be in involution; for details see [45]. Thus, the geodesic motion is completely integrable and all Schouten–Nijenhuis brackets of  ${}^j k$  and  ${}^k l$  must vanish.

## B. Separability of the Klein–Gordon equation

We want to demonstrate that the geometry (2.1) enables us to solve the Klein–Gordon equation by a separation of variables. We will actually prove the separability for all operators based on the Killing tensors  ${}^j k$  and

<sup>3</sup> The off-shell Wahlquist metric can be regained from (2.1) by setting  $Z_\mu = \frac{1}{\beta^2} \sinh^2 \beta x_\mu$ .

<sup>4</sup> Despite the zeros in the denominators, some of the following expressions can be also extended to the  $Z'_\nu = 0$  case because  $S_{\mu,\nu} = -\frac{2Z'_\nu}{Z_\mu - Z_\nu} \left( Z''_\nu - \frac{1}{2} \frac{Z'_\mu{}^2 - Z'_\nu{}^2}{Z_\mu - Z_\nu} \right)$ .

Killing vectors  ${}^j\mathbf{l}$  introduced in (2.11) and (2.12), namely, for operators

$$\mathbf{K}_j = -\nabla_a {}^j\mathbf{k}^{ab} \nabla_b, \quad \mathbf{L}_j = -i {}^j\mathbf{l}^a \nabla_a. \quad (2.13)$$

Operator  $\mathbf{K}_j$  can be rewritten in coordinates using the fact that  $\mathbf{K}_j = -\mathbf{g}^{-\frac{1}{2}} \partial_a \mathbf{g}^{\frac{1}{2}} {}^j\mathbf{k}^{ab} \partial_b$  and employing relation (A7). Here,  $\partial$  denotes the coordinate derivative with respect to the coordinates  $x_\mu, \psi_k$ . The operators  $\mathbf{K}_j$  and  $\mathbf{L}_j$  then read<sup>5</sup>

$$\begin{aligned} \mathbf{K}_j &= \sum_\mu \frac{A_\mu^{(j)}}{U_\mu} \left[ -\frac{\partial}{\partial x_\mu} X_\mu \frac{\partial}{\partial x_\mu} \right. \\ &\quad \left. + \frac{1}{X_\mu} \left[ \sum_k (-Z_\mu)^{N-1-k} \mathbf{L}_k \right]^2 \right], \\ \mathbf{L}_k &= -i \frac{\partial}{\partial \psi_k}. \end{aligned} \quad (2.14)$$

Following the procedure in [13], we show that these operators mutually commute. First, the commutators of  $\mathbf{L}_j$  among themselves and of  $\mathbf{L}_j$  and  $\mathbf{K}_k$  obviously vanish,

$$[\mathbf{L}_j, \mathbf{L}_k] = 0, \quad [\mathbf{L}_j, \mathbf{K}_k] = 0, \quad (2.15)$$

and it remains to prove that

$$[\mathbf{K}_j, \mathbf{K}_k] = 0. \quad (2.16)$$

Operators  $\mathbf{K}_j$  can be written as

$$\mathbf{K}_j = \sum_\mu \frac{A_\mu^{(j)}}{U_\mu} \mathbf{M}_\mu, \quad (2.17)$$

where

$$\mathbf{M}_\mu = -\frac{\partial}{\partial x_\mu} X_\mu \frac{\partial}{\partial x_\mu} + \frac{1}{X_\mu} \left[ \sum_k (-Z_\mu)^{N-1-k} \mathbf{L}_k \right]^2. \quad (2.18)$$

Operator  $\mathbf{M}_\mu$  depends only on a corresponding coordinate  $x_\mu$  and contains a derivative  $\frac{\partial}{\partial x_\mu}$ . Due to this fact, these operators commute,

$$[\mathbf{M}_\mu, \mathbf{M}_\nu] = 0. \quad (2.19)$$

Substituting the inverse relation [cf. identities (A1)]

$$\mathbf{M}_\mu = \sum_j (-Z_\mu)^{N-1-j} \mathbf{K}_j \quad (2.20)$$

into (2.19), we obtain

$$\sum_{j,k} (-Z_\mu)^{N-1-j} (-Z_\nu)^{N-1-k} [\mathbf{K}_j, \mathbf{K}_k] = 0, \quad (2.21)$$

which leads to (2.16).

Therefore, all operators (2.14) mutually commute and, thus, they have a common set of eigenfunctions, which satisfy

$$\mathbf{K}_j \phi = \Xi_j \phi, \quad \mathbf{L}_j \phi = \Psi_j \phi. \quad (2.22)$$

Here,  $\Xi_j$  and  $\Psi_j$  are corresponding eigenvalues which label the eigenfunctions.

By employing the identities (A1), we find the solution in the separated form

$$\phi = \prod_\mu R_\mu \prod_k \exp(i \Psi_k \psi_k), \quad (2.23)$$

where  $R_\mu = R_\mu(x_\mu)$  are single-variable functions which satisfy ordinary differential equations,

$$\left( X_\mu R'_\mu \right)' + \left( \check{\Xi}_\mu - \frac{\check{\Psi}_\mu^2}{X_\mu} \right) R_\mu = 0, \quad (2.24)$$

where

$$\check{\Psi}_\mu = \sum_k \Psi_k (-Z_\mu)^{N-1-k}, \quad \check{\Xi}_\mu = \sum_k \Xi_k (-Z_\mu)^{N-1-k}. \quad (2.25)$$

### C. Solutions of the Einstein equations

In what follows, we find functions  $Z_\mu$  and  $X_\mu$  for which the metric (2.1) solves the vacuum Einstein equations with cosmological constant  $\Lambda$ , which in  $D = 2N$  can be written as

$$\mathbf{Ric} = \frac{\Lambda}{N-1} \mathbf{g}, \quad N > 1. \quad (2.26)$$

We see from (2.7) that we must require

$$S_\mu = S_\nu \quad (2.27)$$

to satisfy the off-diagonal part of Einstein equations. In the four-dimensional case,  $N = 2$ , this requirement says that  $Z''_\mu = 2p$ , i.e.,

$$Z_\mu = px_\mu^2 + q_\mu x_\mu + r_\mu, \quad (2.28)$$

with  $p, q_\mu$ , and  $r_\mu$  being constants. Assumption  $Z'_\mu \neq 0$  requires that  $p \neq 0$  or  $p = 0, q_\mu \neq 0$ .

If  $N > 2$ , then (2.27) implies that functions  $Z_\mu$  meet the condition

$$\frac{Z_\mu'^2 - Z_\nu'^2}{Z_\mu - Z_\nu} = K_\mu + K_\nu + s_{\mu\nu} \quad (2.29)$$

<sup>5</sup> In operator equations we use the convention that the round brackets around a derivative end the action of the derivative to the right, however, the square brackets do not. It means  $[\nabla_a \mathbf{l}^a] = \mathbf{l}^a \nabla_a + (\nabla_a \mathbf{l}^a)$ . Applying the operator on a scalar  $\phi$ , we get  $[\nabla_a \mathbf{l}^a] \phi = (\nabla_a (\mathbf{l}^a \phi)) = \mathbf{l}^a (\nabla_a \phi) + \phi (\nabla_a \mathbf{l}^a)$ .

for the single-variable functions  $K_\mu = K_\mu(x_\mu)$  and constants  $s_{\mu\nu} = s_{\nu\mu}$ . Indeed, if we differentiate (2.27) with respect to  $x_\kappa$ , where  $\kappa \neq \mu, \nu$ , we find out that

$$\left( \frac{Z'_\mu{}^2 - Z'_\kappa{}^2}{Z_\mu - Z_\kappa} \right)_{,\kappa} = L_\kappa \quad (2.30)$$

for the single-variable functions  $L_\kappa = L_\kappa(x_\kappa)$ . Integrating this equation and using the symmetry of the left-hand side, we obtain the condition (2.29).

Next, we multiply the relation (2.29) by  $Z_\mu - Z_\nu$  and differentiate it with respect to both indices, which gives

$$Z'_\mu K'_\nu = Z'_\nu K'_\mu. \quad (2.31)$$

This equation is equivalent to

$$\frac{K'_\mu}{Z'_\mu} = q, \quad (2.32)$$

with  $q$  being a separation constant. By integrating (2.32), we obtain

$$K_\mu = qZ_\mu + p_\mu, \quad (2.33)$$

where  $p_\mu$  are arbitrary constants. Substituting (2.33) into (2.29) yields

$$Z'_\mu{}^2 - qZ_\mu^2 - (p_\mu + p_\nu + s_{\mu\nu})Z_\mu = r_{\mu\nu}, \quad \mu \neq \nu, \quad (2.34)$$

where  $r_{\mu\nu} = r_{\nu\mu}$  are separation constants. Differentiating this relation we find that the expressions  $(p_\mu + p_\nu + s_{\mu\nu})$  and  $r_{\mu\nu}$  do not depend on indices  $\mu, \nu$ . We denote these constants by  $4p$  and  $r$ , respectively, i.e.

$$Z'_\mu{}^2 - qZ_\mu^2 - 4pZ_\mu = r. \quad (2.35)$$

In order to ensure the compatibility with the original equation (2.27), the constant  $q$  must vanish. Thus, the relation (2.35), finally, takes the form

$$Z'_\mu{}^2 - 4pZ_\mu = r. \quad (2.36)$$

Equation (2.36) is satisfied if and only if  $Z_\mu$  has the form (2.28), where  $r_\mu$  are arbitrary constants. However, unlike the  $N = 2$  case, the constants  $q_\mu$  must now satisfy

$$q_\mu^2 - 4pr_\mu = r, \quad (2.37)$$

where  $p \neq 0$  or  $p = 0, r \neq 0$ .

Let us now turn our attention to the trace of the Einstein equations,

$$\mathcal{R} = \frac{2N\Lambda}{N-1}. \quad (2.38)$$

Substituting (2.9) and applying lemma A.1, we obtain

$$-X''_\mu = \frac{2N\Lambda}{N-1}(-Z_\mu)^{N-1} + \sum_{k=0}^{N-2} a_k(-Z_\mu)^k, \quad (2.39)$$

with arbitrary constants  $a_k$ . Equation (2.39) is a simple linear differential equation, where the right-hand side is a polynomial of degree at most  $2N$  in  $x_\mu$ . It has a general solution,

$$X_\mu = \alpha_\mu x_\mu + \beta_\mu + \sum_{k=1}^{N+1} d_k(-Z_\mu)^k, \quad (2.40)$$

where  $\alpha_\mu, \beta_\mu$  are arbitrary coefficients of the homogeneous solution and  $d_k$  correspond to the particular solution. These coefficients are determined by the right-hand side of (2.39) or just replace the arbitrary constants  $a_k$ ,

$$d_{N+1} = \begin{cases} 0 & p \neq 0 \\ -\frac{2}{N^2-1} \frac{\Lambda}{r} & p = 0, r \neq 0 \end{cases}, \quad d_N = \begin{cases} \frac{1}{(N-1)(2N-1)} \frac{\Lambda}{p} & p \neq 0 \\ \text{const.} \in \mathbb{R} & p = 0, r \neq 0 \end{cases}, \quad (2.41)$$

$$d_{N-1} = \text{const.} \in \mathbb{R},$$

$$\vdots$$

$$d_1 = \text{const.} \in \mathbb{R}.$$

Substituting (2.40) into the full Einstein equations and employing (A1) and (A2), we find

$$\alpha_\mu q_\mu - 2\beta_\mu p = s, \quad (2.42)$$

where  $s$  is a constant.

Thus, the full Einstein equations are satisfied if and only if the functions  $Z_\mu$  are given by (2.28), (2.37), and the functions  $X_\mu$  by (2.40)–(2.42).

These relations are also valid for  $N = 2$ , because the condition (2.36) is met here anyway; however, here it is enforced by the diagonal part instead of the off-diagonal part of the Einstein equations.

For  $p \neq 0$ , we can introduce the new coordinates

$$\check{x}_\mu = \sqrt{p} \left( x_\mu + \frac{q_\mu}{2p} \right), \quad \check{\psi}_j = \frac{1}{\sqrt{p}} \sum_{k=0}^{N-1-j} \frac{(j+k)!}{j!} \left( -\frac{r}{4p} \right)^k \psi_{j+k}. \quad (2.43)$$

In these coordinates, the metric (2.1) takes the form

$$g = \sum_\mu \left[ \frac{\check{U}_\mu}{\check{X}_\mu} (\mathbf{d}\check{x}_\mu)^2 + \frac{\check{X}_\mu}{\check{U}_\mu} \left( \sum_j \check{A}_\mu^{(j)} \mathbf{d}\check{\psi}_j \right)^2 \right], \quad (2.44)$$

where

$$\check{Z}_\mu = \check{x}_\mu^2, \quad \check{X}_\mu = \sum_{k=0}^N \check{c}_k (-\check{x}_\mu^2)^k + \check{b}_\mu \check{x}_\mu. \quad (2.45)$$

Here, the coefficient of the highest power of  $x_\mu$  is

$$\check{c}_N = \frac{\Lambda}{(N-1)(2N-1)}, \quad (2.46)$$



and the remaining coefficients  $\check{c}_k$ ,  $\check{b}_\mu$  are arbitrary constants replacing the  $d$ 's,  $\alpha$ 's, and  $\beta$ 's. This is a standard form of the Kerr–NUT–(A)dS spacetime [9], and the 2-form (2.10) is a standard principal conformal Killing–Yano tensor. The metric is written in a Euclidean form, however, for a suitable choice of ranges of coordinates, it can be Wick-rotated to a Lorentzian signature (see [9, 42]).

For  $p = 0$  and  $r \neq 0$ , we define other coordinates<sup>6</sup>

$$\check{x}_\mu = \pm_\mu \sqrt{r} x_\mu + r_\mu, \quad \check{\psi}_j = \frac{1}{\sqrt{r}} \psi_j. \quad (2.47)$$

In these coordinates, the metric (2.1) takes the form (2.44), where

$$\check{Z}_\mu = \check{x}_\mu, \quad \check{X}_\mu = \sum_{k=1}^{N+1} \check{c}_k (-\check{x}_\mu)^k + \check{b}_\mu. \quad (2.48)$$

Here, the coefficient of the highest power of  $x_\mu$  is

$$\check{c}_{N+1} = -\frac{2\Lambda}{N^2 - 1}, \quad (2.49)$$

and the remaining coefficients  $\check{c}_k$  and  $\check{b}_\mu$  are again arbitrary constants replacing the  $d$ 's,  $\alpha$ 's, and  $\beta$ 's.

It can be verified that the metric (2.48) is Kähler and the 2-form

$$\Omega = \sum_\mu e^\mu \wedge \hat{e}^\mu \quad (2.50)$$

is the corresponding Kähler 2-form. This Einstein–Kähler metric can be also obtained directly from the even-dimensional Kerr–NUT–(A)dS metric using a particular scaling limit [46] and it is often used to construct a one-dimensional higher Sasakian metric.

Another approach of constructing these metrics is by employing an analog of the Euclidean BPS limit. Even and odd-dimensional Kerr–NUT–(A)dS spacetime then leads to Ricci-flat–Kähler and Einstein–Sasakian metrics, respectively, see [9, 10].

Unfortunately, the metric (2.48) has a Euclidean signature and cannot be simply Wick-rotated to a Lorentzian regime. This is why it was not much studied in  $D = 4$ .

For further reference, we briefly discuss the  $N = 1$  case, when the metric (2.1) takes the form

$$g = \frac{1}{X_1} (\mathbf{d}x_1)^2 + X_1 (\mathbf{d}\psi_0)^2. \quad (2.51)$$

The corresponding Ricci tensor and scalar curvature are

$$\mathbf{Ric} = -\frac{1}{2} X_1'' g, \quad \mathcal{R} = -X_1''. \quad (2.52)$$

The closest two-dimensional analog of the Einstein equations reads  $\mathcal{R} - \Lambda = 0$ , see [47]. This equation is met if and only if

$$X_1 = \frac{\Lambda}{2} (-x_1^2) + b_1 x_1 + c_0, \quad (2.53)$$

where  $b_1$  and  $c_0$  are arbitrary constants.

### III. WARPED KLEIN–GORDON SEPARABLE SPACETIMES

In the previous section we introduced the Klein–Gordon simple separable metric (2.1) and found two important solutions of the Einstein equations, (2.45) and (2.48). Although the former is the Kerr–NUT–(A)dS spacetime, which has a clear physical significance, the latter cannot be even simply Wick-rotated to a Lorentzian signature. However, in higher dimensions, it could be used to construct physically interesting warped metrics. In addition, due to the simplicity of warped geometries, such metrics also could be Klein–Gordon separable.

#### A. Warped geometry

Consider the direct product  $M = \tilde{M} \times \bar{M}$  of two manifolds with dimensions  $\tilde{D}$  and  $\bar{D}$ , respectively. The warped space is the manifold  $M$  of dimension  $D = \tilde{D} + \bar{D}$  equipped with the metric

$$g = \tilde{g} + \tilde{w}^2 \bar{g}, \quad (3.1)$$

where  $\tilde{g}$ ,  $\bar{g}$  are metrics on the manifolds  $\tilde{M}$ ,  $\bar{M}$ , respectively, and  $\tilde{w}^2$  is a warped factor. We assume that the tilded and barred objects depend only on a position in  $\tilde{M}$  or  $\bar{M}$ , and they are also nontrivial only in tilded or barred directions.

We define torsion-free covariant derivatives  $\nabla$ ,  $\tilde{\nabla}$ , and  $\bar{\nabla}$  satisfying  $\nabla g = 0$ ,  $\tilde{\nabla} \tilde{g} = 0$ , and  $\bar{\nabla} \bar{g} = 0$ , respectively. The Christoffel coefficients with respect to the adjusted coordinates then satisfy the relation

$$\Gamma_{ab}^c = \tilde{\Gamma}_{ab}^c + \bar{\Gamma}_{ab}^c + \Lambda_{ab}^c, \quad (3.2)$$

where

$$\Lambda_{ab}^c = 2\tilde{\lambda}_{(a} \bar{\delta}_{b)}^c - \tilde{w}^2 \tilde{\lambda}^c \bar{g}_{ab} \quad (3.3)$$

with the 1-form  $\tilde{\lambda}$  being a logarithmic derivative of a warped factor,

$$\tilde{\lambda} = \mathbf{d} \ln \tilde{w}, \quad \tilde{\lambda}^2 = \tilde{g}^{ab} \tilde{\lambda}_a \tilde{\lambda}_b. \quad (3.4)$$

The Riemann tensor, Ricci tensor, and scalar curvature read

$$\begin{aligned} R_{ab}{}^c{}_d &= \tilde{R}_{ab}{}^c{}_d + \frac{2}{\tilde{w}} \tilde{H}_{d[a} \bar{\delta}_{b]}^c - 2\tilde{w} \tilde{g}^{ce} \tilde{H}_{e[a} \bar{g}_{b]d} \\ &\quad + \bar{R}_{ab}{}^c{}_d - 2\tilde{w}^2 \tilde{\lambda}^2 \bar{\delta}_{[a}^c \bar{g}_{b]d}, \\ \mathbf{Ric}_{ab} &= \tilde{\mathbf{Ric}}_{ab} - \frac{\bar{D}}{\tilde{w}} \tilde{H}_{ab} \\ &\quad + \bar{\mathbf{Ric}}_{ab} - \tilde{w}^2 \left( \frac{\tilde{H}}{\tilde{w}} + (\bar{D} - 1) \tilde{\lambda}^2 \right) \bar{g}_{ab}, \\ \mathcal{R} &= \tilde{\mathcal{R}} + \frac{\bar{\mathcal{R}}}{\tilde{w}^2} - \frac{2\bar{D}}{\tilde{w}} \tilde{H} - \bar{D}(\bar{D} - 1) \tilde{\lambda}^2. \end{aligned} \quad (3.5)$$

<sup>6</sup> Symbol  $\pm_\mu$  denotes the sign which depends on index  $\mu$ .

Here, the Hessian tensor  $\tilde{\mathbf{H}}$  and scalar  $\tilde{\mathcal{H}}$  are

$$\tilde{\mathbf{H}} = \tilde{\nabla} \tilde{\nabla} \tilde{w}, \quad \tilde{\mathcal{H}} = \tilde{g}^{ab} \tilde{\mathbf{H}}_{ab}. \quad (3.6)$$

The vacuum Einstein equations  $\mathbf{Ric} = \frac{2\Lambda}{D-2} \mathbf{g}$  split into two parts,

$$\tilde{\mathbf{Ric}} = \frac{\bar{D}}{\tilde{w}} \tilde{\mathbf{H}} + \frac{2\Lambda}{D-2} \tilde{g} \quad (3.7)$$

and

$$\bar{\mathbf{Ric}} = \tilde{w}^2 \left( \frac{\tilde{\mathcal{H}}}{\tilde{w}} + (\bar{D} - 1) \tilde{\lambda}^2 + \frac{2\Lambda}{D-2} \right) \tilde{g}. \quad (3.8)$$

However, the right-hand side of (3.8) should not depend on the position in  $\tilde{M}$ , so the coefficient in front of  $\tilde{g}$  must be constant. We denote this constant by  $\Upsilon$ , i.e.,

$$\Upsilon = \tilde{w}^2 \left( \frac{\tilde{\mathcal{H}}}{\tilde{w}} + (\bar{D} - 1) \tilde{\lambda}^2 + \frac{2\Lambda}{D-2} \right). \quad (3.9)$$

Assume that  $\tilde{\mathbf{k}}$  and  $\bar{\mathbf{k}}$  are rank-2 Killing tensors of the metrics  $\tilde{g}$  and  $\bar{g}$ , respectively. Then it was shown in [43] that

$$\tilde{\mathbf{k}} + \frac{\tilde{A}}{\tilde{w}^2} \tilde{g}^{-1} \quad \text{and} \quad \bar{\mathbf{k}} \quad (3.10)$$

are Killing tensors of the metric  $\mathbf{g}$ , provided that  $\tilde{A}$  is a function on  $\tilde{M}$  satisfying

$$\tilde{\nabla}^a \tilde{A} = 2(\tilde{A} \tilde{g}^{ab} - \tilde{\mathbf{k}}^{ab}) \tilde{\lambda}_b. \quad (3.11)$$

### B. Off-shell metric

Consider a warped metric (3.1), where both  $\tilde{g}$  and  $\bar{g}$  are the off-shell metrics (2.1) of dimensions  $\bar{D} = 2\tilde{N}$  and  $\bar{D} = 2\bar{N}$ , respectively, and the warped function is given by<sup>7</sup>

$$\tilde{w}^2 = \tilde{A}^{(\tilde{N})} = \tilde{Z}_1 \dots \tilde{Z}_{\tilde{N}}. \quad (3.12)$$

The determinant  $\mathbf{g}$  of this metric in the coordinates  $\tilde{x}_{\tilde{\mu}}$ ,  $\tilde{\psi}_{\tilde{k}}$ ,  $\tilde{x}_{\bar{\mu}}$ ,  $\tilde{\psi}_{\bar{k}}$  reads

$$\mathbf{g} = \tilde{w}^{2\tilde{N}} \tilde{U}^2 \bar{U}^2. \quad (3.13)$$

The 1-form (3.4) is given by

$$\tilde{\lambda} = \frac{1}{2} \sum_{\tilde{\mu}} \frac{\tilde{Z}'_{\tilde{\mu}}}{\tilde{Z}_{\tilde{\mu}}} d\tilde{x}_{\tilde{\mu}}, \quad \tilde{\lambda}^2 = \frac{1}{4} \sum_{\tilde{\mu}} \frac{\tilde{Z}'_{\tilde{\mu}}{}^2}{\tilde{Z}_{\tilde{\mu}}^2} \frac{\tilde{X}_{\tilde{\mu}}}{\tilde{U}_{\tilde{\mu}}}. \quad (3.14)$$

The Hessian tensor and scalar (3.6) read

$$\begin{aligned} \tilde{\mathbf{H}} &= \frac{1}{4} \sum_{\tilde{\mu}, \tilde{\nu}} \frac{\tilde{w}}{\tilde{Z}_{\tilde{\mu}}} \left[ \left( \frac{\tilde{Z}'_{\tilde{\nu}}{}^2}{\tilde{Z}_{\tilde{\nu}}} \frac{\tilde{X}_{\tilde{\nu}}}{\tilde{U}_{\tilde{\nu}}} \right)_{,\tilde{\mu}} \tilde{e}^{\tilde{\mu}} \tilde{e}^{\tilde{\nu}} + \frac{\tilde{Z}'_{\tilde{\nu}}{}^2}{\tilde{Z}_{\tilde{\nu}}} \left( \frac{\tilde{X}_{\tilde{\nu}}}{\tilde{U}_{\tilde{\nu}}} \right)_{,\tilde{\mu}} \tilde{e}^{\tilde{\mu}} \tilde{e}^{\tilde{\nu}} \right] \\ &\quad + \sum_{\substack{\tilde{\mu}, \tilde{\nu} \\ \tilde{\nu} \neq \tilde{\mu}}} \frac{\tilde{w}}{\tilde{Z}_{\tilde{\mu}} - \tilde{Z}_{\tilde{\nu}}} \left( \frac{\tilde{Z}'_{\tilde{\nu}}{}^2}{\tilde{Z}_{\tilde{\nu}}} - \frac{\tilde{Z}'_{\tilde{\mu}}{}^2}{\tilde{Z}_{\tilde{\mu}}} \right) \sqrt{\frac{\tilde{X}_{\tilde{\mu}}}{\tilde{U}_{\tilde{\mu}}} \frac{\tilde{X}_{\tilde{\nu}}}{\tilde{U}_{\tilde{\nu}}}} \tilde{e}^{\tilde{\mu}} \tilde{e}^{\tilde{\nu}}, \\ \tilde{\mathcal{H}} &= \frac{1}{2} \sum_{\tilde{\mu}} \frac{\tilde{w}}{\sqrt{\tilde{Z}_{\tilde{\mu}}}} \left( \frac{\tilde{Z}'_{\tilde{\mu}}}{\sqrt{\tilde{Z}_{\tilde{\mu}}}} \tilde{X}_{\tilde{\mu}} \right)' \frac{1}{\tilde{U}_{\tilde{\mu}}}. \end{aligned} \quad (3.15)$$

The Riemann tensor, Ricci tensor, and scalar curvature can be then calculated directly by substituting (B4), (2.7), (2.9), (3.15), and (3.14) into the general expressions (3.5).

Employing (3.10), (3.11), and (A3), it can be easily verified that

$${}^i \mathbf{k} = {}^i \tilde{\mathbf{k}} + \frac{\tilde{A}^{(i)}}{\tilde{w}^2} \tilde{g}^{-1}, \quad {}^{\tilde{N}+i} \mathbf{k} = {}^i \bar{\mathbf{k}} \quad (3.16)$$

are rank-2 Killing tensors of the warped metric. Because the warped function  $\tilde{w}$  does not depend on the coordinates  $\tilde{\psi}_{\tilde{j}}$ , the Killing vectors remain unchanged, i.e.,

$${}^i \mathbf{l} = {}^i \tilde{\mathbf{l}}, \quad {}^{\tilde{N}+i} \mathbf{l} = {}^i \bar{\mathbf{l}}. \quad (3.17)$$

Again, since we will show that all operators (3.18) constructed below mutually commute, the corresponding phase-space observables  ${}^j \mathbf{k}^{ab} \mathbf{p}_a \mathbf{p}_b$ ,  ${}^k \mathbf{l}^a \mathbf{p}_a$  must be in involution. Thus, the geodesic motion is completely integrable and all Schouten–Nijenhuis brackets of  ${}^j \mathbf{k}$  and  ${}^k \mathbf{l}$  must vanish.

### C. Separability of the Klein–Gordon equation

We again define operators

$$\mathbf{K}_j = -\nabla_a {}^j \mathbf{k}^{ab} \nabla_b, \quad \mathbf{L}_j = -i {}^j \mathbf{l}^a \nabla_a, \quad (3.18)$$

based on Killing tensors  ${}^j \mathbf{k}$  (3.16) and Killing vectors  ${}^j \mathbf{l}$  (3.17). Operators (3.18) can be also expressed in coordinates  $\tilde{x}_{\tilde{\mu}}$ ,  $\tilde{\psi}_{\tilde{k}}$ ,  $\tilde{x}_{\bar{\mu}}$ ,  $\tilde{\psi}_{\bar{k}}$  using  $\mathbf{K}_j = -\mathbf{g}^{-\frac{1}{2}} \partial_a \mathbf{g}^{\frac{1}{2}} {}^j \mathbf{k}^{ab} \partial_b$  and

<sup>7</sup> This form ensures the separability of the Klein–Gordon equation (3.27) below. It is probably not the most general ansatz with this property; the question of the most general warp factor admitting the separability remains open.

employing relation (A7),

$$\begin{aligned}
K_{\tilde{j}} &= \sum_{\tilde{\mu}} \frac{\tilde{A}_{\tilde{\mu}}^{(\tilde{j})}}{\tilde{U}_{\tilde{\mu}}} \left[ -\frac{1}{\tilde{Z}_{\tilde{\mu}}^{\frac{\tilde{N}}{2}}} \frac{\partial}{\partial \tilde{x}_{\tilde{\mu}}} \tilde{Z}_{\tilde{\mu}}^{\frac{\tilde{N}}{2}} \tilde{X}_{\tilde{\mu}} \frac{\partial}{\partial \tilde{x}_{\tilde{\mu}}} \right. \\
&\quad \left. + \frac{1}{\tilde{X}_{\tilde{\mu}}} \left[ \sum_{\tilde{k}} (-\tilde{Z}_{\tilde{\mu}})^{\tilde{N}-1-\tilde{k}} \mathbf{L}_{\tilde{k}} \right]^2 \right] + \frac{\tilde{A}^{(\tilde{j})}}{\tilde{A}^{(\tilde{N})}} K_{\tilde{N}}, \\
K_{\tilde{N}+\tilde{j}} &= \sum_{\tilde{\mu}} \frac{\tilde{A}_{\tilde{\mu}}^{(\tilde{j})}}{\tilde{U}_{\tilde{\mu}}} \left[ -\frac{\partial}{\partial \tilde{x}_{\tilde{\mu}}} \tilde{X}_{\tilde{\mu}} \frac{\partial}{\partial \tilde{x}_{\tilde{\mu}}} \right. \\
&\quad \left. + \frac{1}{\tilde{X}_{\tilde{\mu}}} \left[ \sum_{\tilde{k}} (-\tilde{Z}_{\tilde{\mu}})^{\tilde{N}-1-\tilde{k}} \mathbf{L}_{\tilde{N}+\tilde{k}} \right]^2 \right], \\
\mathbf{L}_{\tilde{k}} &= -i \frac{\partial}{\partial \psi_{\tilde{k}}} , \quad \mathbf{L}_{\tilde{N}+\tilde{k}} = -i \frac{\partial}{\partial \psi_{\tilde{k}}} .
\end{aligned} \tag{3.19}$$

It can be shown that these operators mutually commute. Indeed, commutators of  $\mathbf{L}_j$  among themselves and commutators of  $\mathbf{L}_j$  and  $\mathbf{K}_k$  vanish,

$$[\mathbf{L}_j, \mathbf{L}_k] = 0, \quad [\mathbf{L}_j, \mathbf{K}_k] = 0. \tag{3.20}$$

Operators  $\mathbf{K}_j$  have the form

$$K_{\tilde{j}} = \sum_{\tilde{\mu}} \frac{\tilde{A}_{\tilde{\mu}}^{(\tilde{j})}}{\tilde{U}_{\tilde{\mu}}} \tilde{\mathbf{M}}_{\tilde{\mu}} + \frac{\tilde{A}^{(\tilde{j})}}{\tilde{A}^{(\tilde{N})}} K_{\tilde{N}}, \quad K_{\tilde{N}+\tilde{j}} = \sum_{\tilde{\mu}} \frac{\tilde{A}_{\tilde{\mu}}^{(\tilde{j})}}{\tilde{U}_{\tilde{\mu}}} \tilde{\mathbf{M}}_{\tilde{\mu}}, \tag{3.21}$$

where

$$\begin{aligned}
\tilde{\mathbf{M}}_{\tilde{\mu}} &= -\frac{1}{\tilde{Z}_{\tilde{\mu}}^{\frac{\tilde{N}}{2}}} \frac{\partial}{\partial \tilde{x}_{\tilde{\mu}}} \tilde{Z}_{\tilde{\mu}}^{\frac{\tilde{N}}{2}} \tilde{X}_{\tilde{\mu}} \frac{\partial}{\partial \tilde{x}_{\tilde{\mu}}} \\
&\quad + \frac{1}{\tilde{X}_{\tilde{\mu}}} \left[ \sum_{\tilde{k}} (-\tilde{Z}_{\tilde{\mu}})^{\tilde{N}-1-\tilde{k}} \mathbf{L}_{\tilde{k}} \right]^2, \\
\tilde{\mathbf{M}}_{\tilde{\mu}} &= -\frac{\partial}{\partial \tilde{x}_{\tilde{\mu}}} \tilde{X}_{\tilde{\mu}} \frac{\partial}{\partial \tilde{x}_{\tilde{\mu}}} + \frac{1}{\tilde{X}_{\tilde{\mu}}} \left[ \sum_{\tilde{k}} (-\tilde{Z}_{\tilde{\mu}})^{\tilde{N}-1-\tilde{k}} \mathbf{L}_{\tilde{N}+\tilde{k}} \right]^2.
\end{aligned} \tag{3.22}$$

Commutators of  $\tilde{\mathbf{M}}_{\tilde{\mu}}$  and  $\tilde{\mathbf{M}}_{\tilde{\nu}}$  among themselves vanish,

$$[\tilde{\mathbf{M}}_{\tilde{\mu}}, \tilde{\mathbf{M}}_{\tilde{\nu}}] = 0, \quad [\tilde{\mathbf{M}}_{\tilde{\mu}}, \tilde{\mathbf{M}}_{\tilde{\nu}}] = 0, \tag{3.23}$$

which, much like in Sec. II B, implies

$$[K_{\tilde{j}}, K_{\tilde{k}}] = 0, \quad [K_{\tilde{N}+\tilde{j}}, K_{\tilde{N}+\tilde{k}}] = 0. \tag{3.24}$$

In the first equation we have also used the relation

$$\left[ \sum_{\tilde{\mu}} \frac{\tilde{A}_{\tilde{\mu}}^{(\tilde{j})}}{\tilde{U}_{\tilde{\mu}}} \tilde{\mathbf{M}}_{\tilde{\mu}}, \frac{\tilde{A}^{(\tilde{k})}}{\tilde{A}^{(\tilde{N})}} K_{\tilde{N}} \right] = \left[ \sum_{\tilde{\mu}} \frac{\tilde{A}_{\tilde{\mu}}^{(\tilde{k})}}{\tilde{U}_{\tilde{\mu}}} \tilde{\mathbf{M}}_{\tilde{\mu}}, \frac{\tilde{A}^{(\tilde{j})}}{\tilde{A}^{(\tilde{N})}} K_{\tilde{N}} \right], \tag{3.25}$$

which is a consequence of identity (A8). Finally, using the second equation in (3.24), we can also prove the relation

$$[K_{\tilde{j}}, K_{\tilde{N}+\tilde{k}}] = 0. \tag{3.26}$$

We have thus proved that operators (3.19) mutually commute and, therefore, they have a common set of eigenfunctions; i.e., they satisfy

$$K_j \phi = \Xi_j \phi, \quad \mathbf{L}_j \phi = \Psi_j \phi, \tag{3.27}$$

where  $\Xi_j$  and  $\Psi_j$  are corresponding eigenvalues.

Employing identities (A1) and (A4), we find the solution in the separated form,

$$\phi = \prod_{\tilde{\mu}} \tilde{R}_{\tilde{\mu}} \prod_{\tilde{\mu}} \bar{R}_{\tilde{\mu}} \prod_{\tilde{k}} \exp(i \Psi_{\tilde{k}} \tilde{\psi}_{\tilde{k}}) \prod_{\tilde{k}} \exp(i \Psi_{\tilde{N}+\tilde{k}} \bar{\psi}_{\tilde{k}}), \tag{3.28}$$

where  $\tilde{R}_{\tilde{\mu}} = \tilde{R}_{\tilde{\mu}}(\tilde{x}_{\tilde{\mu}})$  and  $\bar{R}_{\tilde{\mu}} = \bar{R}_{\tilde{\mu}}(\bar{x}_{\tilde{\mu}})$  are single-variable functions which satisfy ordinary differential equations,

$$\begin{aligned}
\left( \tilde{Z}_{\tilde{\mu}}^{\frac{\tilde{N}}{2}} \tilde{X}_{\tilde{\mu}} \tilde{R}'_{\tilde{\mu}} \right)' + \left( \check{\Xi}_{\tilde{\mu}} - \frac{\check{\Psi}_{\tilde{\mu}}^2}{\tilde{X}_{\tilde{\mu}}} + \frac{\Xi_{\tilde{N}}}{\tilde{Z}_{\tilde{\mu}}} \right) \tilde{Z}_{\tilde{\mu}}^{\frac{\tilde{N}}{2}} \tilde{R}_{\tilde{\mu}} &= 0, \\
\left( \bar{X}_{\tilde{\mu}} \bar{R}'_{\tilde{\mu}} \right)' + \left( \check{\Xi}_{\tilde{\mu}} - \frac{\check{\Psi}_{\tilde{\mu}}^2}{\bar{X}_{\tilde{\mu}}} \right) \bar{R}_{\tilde{\mu}} &= 0,
\end{aligned} \tag{3.29}$$

where

$$\begin{aligned}
\check{\Xi}_{\tilde{\mu}} &= \sum_{\tilde{k}} \Xi_{\tilde{k}} (-\tilde{Z}_{\tilde{\mu}})^{\tilde{N}-1-\tilde{k}}, \\
\check{\Psi}_{\tilde{\mu}} &= \sum_{\tilde{k}} \Psi_{\tilde{k}} (-\tilde{Z}_{\tilde{\mu}})^{\tilde{N}-1-\tilde{k}}, \\
\check{\Xi}_{\tilde{\mu}} &= \sum_{\tilde{k}} \Xi_{\tilde{N}+\tilde{k}} (-\bar{Z}_{\tilde{\mu}})^{\tilde{N}-1-\tilde{k}}, \\
\check{\Psi}_{\tilde{\mu}} &= \sum_{\tilde{k}} \Psi_{\tilde{N}+\tilde{k}} (-\bar{Z}_{\tilde{\mu}})^{\tilde{N}-1-\tilde{k}}.
\end{aligned} \tag{3.30}$$

#### D. Solutions of the Einstein equations

In the following, we find functions  $\tilde{Z}_{\tilde{\mu}}$ ,  $\tilde{X}_{\tilde{\mu}}$ ,  $\bar{Z}_{\tilde{\mu}}$ , and  $\bar{X}_{\tilde{\mu}}$  for which the warped metric solves the vacuum Einstein equations.

We see from the off-diagonal part of (3.7) that

$$\tilde{S}_{\tilde{\mu}} + \bar{N} \frac{\tilde{Z}_{\tilde{\mu}}'^2}{\tilde{Z}_{\tilde{\mu}}} = \tilde{S}_{\tilde{\nu}} + \bar{N} \frac{\tilde{Z}_{\tilde{\nu}}'^2}{\tilde{Z}_{\tilde{\nu}}}. \tag{3.31}$$

In the  $\tilde{N} = 2$  case, this condition says that

$$\tilde{Z}_{\tilde{\mu}}'' + \bar{N} \frac{\tilde{Z}_{\tilde{\mu}}'^2}{\tilde{Z}_{\tilde{\mu}}} = 2\tilde{p}(2\bar{N} + 1), \tag{3.32}$$

where  $\tilde{p}$  is an arbitrary constant.

If  $\tilde{N} > 2$ , then (3.31) implies

$$\tilde{Z}_{\tilde{\mu}}'^2 - \tilde{q} \tilde{Z}_{\tilde{\mu}}^2 - 4\tilde{p} \tilde{Z}_{\tilde{\mu}} = \tilde{r} \tag{3.33}$$

similar to the relation between (2.27) and (2.35). However, in order to satisfy (3.31), the constants  $\tilde{q}$  and  $\tilde{r}$  must vanish. Thus, the relation (3.33) takes the form

$$\tilde{Z}_{\tilde{\mu}}'^2 - 4\tilde{p} \tilde{Z}_{\tilde{\mu}} = 0, \tag{3.34}$$



with  $\tilde{p}$  being a nonzero constant. Equation (3.34) is satisfied if and only if

$$\tilde{Z}_{\tilde{\mu}} = \tilde{p}\tilde{x}_{\tilde{\mu}}^2 + \tilde{q}_{\tilde{\mu}}\tilde{x}_{\tilde{\mu}} + \tilde{r}_{\tilde{\mu}}, \quad (3.35)$$

where

$$\tilde{q}_{\tilde{\mu}}^2 - 4\tilde{p}\tilde{r}_{\tilde{\mu}} = 0. \quad (3.36)$$

In the  $\tilde{N} = 1, 2$  cases, the form of (3.35) with the condition (3.36) is enforced by the diagonal part of (3.7). In particular, in the  $\tilde{N} = 1$  case, it simply implies condition (3.34). For  $\tilde{N} = 2$ , it gives

$$\left( \tilde{N} \frac{\tilde{Z}_{\tilde{\mu}}'^2}{\tilde{Z}_{\tilde{\mu}}} + \frac{\tilde{Z}_{\tilde{\mu}}'^2 - \tilde{Z}_{\tilde{\nu}}'^2}{\tilde{Z}_{\tilde{\mu}} - \tilde{Z}_{\tilde{\nu}}} \right)_{,\tilde{\mu}} = 0. \quad (3.37)$$

Integrating this equation, multiplying by  $\tilde{Z}_{\tilde{\mu}} - \tilde{Z}_{\tilde{\nu}}$ , and differentiating with respect to  $\tilde{x}_{\tilde{\mu}}$  and  $\tilde{x}_{\tilde{\nu}}$ , we find that

$$\frac{1}{\tilde{Z}_{\tilde{\mu}}} \left( \frac{\tilde{Z}_{\tilde{\mu}}'^2}{\tilde{Z}_{\tilde{\mu}}} \right)' = \tilde{t}_{\tilde{\mu}}, \quad (3.38)$$

where  $\tilde{t}_{\tilde{\mu}}$  is a separation constant. This equation leads to

$$\tilde{Z}_{\tilde{\mu}}'^2 = \tilde{t}_{\tilde{\mu}}\tilde{Z}_{\tilde{\mu}}^2 + \tilde{u}_{\tilde{\mu}}\tilde{Z}_{\tilde{\mu}}, \quad \tilde{Z}_{\tilde{\mu}}'' = \tilde{t}_{\tilde{\mu}}\tilde{Z}_{\tilde{\mu}} + \frac{1}{2}\tilde{u}_{\tilde{\mu}}, \quad (3.39)$$

with  $\tilde{u}_{\tilde{\mu}}$  being an arbitrary constant. Substituting (3.39) into (3.32), we determine the constants  $\tilde{t}_{\tilde{\mu}} = 0$  and  $\tilde{u}_{\tilde{\mu}} = 4\tilde{p}$  which yields (3.34).

Since  $\tilde{Z}_{\tilde{\mu}}$  is a quadratic function (3.35) with  $\tilde{p} \neq 0$ , we can employ the coordinate freedom to transform the tilded coordinates using transformation (2.43) and drop the caron sign. This transformation preserves the form of the orthonormal frame, and it effectively sets the functions  $\tilde{Z}_{\tilde{\mu}}$  to  $\tilde{x}_{\tilde{\mu}}^2$ . Equation (3.9) can then be rewritten as

$$\sum_{\tilde{\mu}} \frac{1}{\tilde{U}_{\tilde{\mu}}} \left[ \frac{\tilde{X}_{\tilde{\mu}}'}{\tilde{x}_{\tilde{\mu}}} + (2\tilde{N} - 1) \frac{\tilde{X}_{\tilde{\mu}}}{\tilde{x}_{\tilde{\mu}}^2} - \frac{\Upsilon}{\tilde{x}_{\tilde{\mu}}^2} + \frac{\Lambda}{\tilde{N} - 1} (-\tilde{x}_{\tilde{\mu}}^2)^{\tilde{N} - 1} \right] = 0. \quad (3.40)$$

Substituting (2.9) and applying lemma A.1, we obtain

$$\tilde{x}_{\tilde{\mu}} \tilde{X}_{\tilde{\mu}}' + (2\tilde{N} - 1) \tilde{X}_{\tilde{\mu}} = \Upsilon + \sum_{\tilde{k}=0}^{\tilde{N}-2} \tilde{a}_{\tilde{k}} \tilde{x}_{\tilde{\mu}}^{2(\tilde{k}+1)} + \frac{\Lambda}{\tilde{N} - 1} (-\tilde{x}_{\tilde{\mu}}^2)^{\tilde{N}}. \quad (3.41)$$

Equation (3.41) is a simple linear differential equation, where the right-hand side is a polynomial of degree  $2\tilde{N}$  in  $\tilde{x}_{\tilde{\mu}}$ . It has a general solution,

$$\tilde{X}_{\tilde{\mu}} = \sum_{\tilde{k}=0}^{\tilde{N}} \tilde{c}_{\tilde{k}} (-\tilde{x}_{\tilde{\mu}}^2)^{\tilde{k}} + \frac{\tilde{b}_{\tilde{\mu}}}{\tilde{x}_{\tilde{\mu}}^{2\tilde{N}-1}}, \quad (3.42)$$

where

$$\tilde{c}_{\tilde{N}} = \frac{\Lambda}{(2\tilde{N} - 1)(\tilde{N} - 1)}, \quad \tilde{c}_0 = \frac{\Upsilon}{2\tilde{N} - 1}, \quad (3.43)$$

and coefficients  $\tilde{b}_{\tilde{\mu}}, \tilde{c}_{\tilde{k}}, \tilde{k} \in 1, \dots, \tilde{N} - 1$  are arbitrary constants. We thus found metric functions for which the proportionality factor (3.9) is a constant. Surprisingly, employing identities (A1), the tilded part (3.7) of the Einstein equations is already satisfied.

The barred part (3.8) of the Einstein equations leads to the solutions (2.45) or (2.48) with the cosmological constant  $\bar{\Lambda} = (\tilde{N} - 1)\Upsilon$ , i.e.,

$$\bar{X}_{\bar{\mu}} = \sum_{\bar{k}=0}^{\bar{N}} \bar{c}_{\bar{k}} (-\bar{x}_{\bar{\mu}}^2)^{\bar{k}} + \bar{b}_{\bar{\mu}} \bar{x}_{\bar{\mu}}, \quad \bar{c}_{\bar{N}} = \frac{\Upsilon}{2\bar{N} - 1}, \quad (3.44)$$

where  $\bar{b}_{\bar{\mu}}, \bar{c}_{\bar{k}}, \bar{k} \in 0, \dots, \bar{N} - 1$  are arbitrary constants, or

$$\bar{X}_{\bar{\mu}} = \sum_{\bar{k}=1}^{\bar{N}+1} \bar{c}_{\bar{k}} (-\bar{x}_{\bar{\mu}}^2)^{\bar{k}} + \bar{b}_{\bar{\mu}}, \quad \bar{c}_{\bar{N}+1} = -\frac{2\Upsilon}{\bar{N} + 1}, \quad (3.45)$$

where  $\bar{b}_{\bar{\mu}}, \bar{c}_{\bar{k}}, \bar{k} \in 1, \dots, \bar{N}$  are arbitrary constants.

The warped solution with barred functions (3.44) have been recently obtained in [42] and, as discussed there, it also belongs to class of generalized Kerr–NUT–(A)dS spacetimes [38, 39]. The warped metric with barred functions (3.45) constitutes new solutions of the Einstein equations.

#### IV. CONCLUSIONS

In this paper we have generalized Carter's ansatz for the metric leading to a separable Klein–Gordon equation to higher dimensions and thus defined the class of Klein–Gordon simple separable metrics. For these metrics, we have solved the Klein–Gordon equation by separation of variables, employing the symmetries encoded in Killing vectors and rank-two Killing tensors. The class of such metrics enlarges previously studied off-shell Kerr–NUT–(A)dS spaces. Solving the Einstein equations in this class, we have regained the on-shell Kerr–NUT–(A)dS spacetimes and the Euclidean Einstein–Kähler metrics which, as demonstrated in [9, 10, 46], can be obtained as a particular limit of the Kerr–NUT–(A)dS spaces as well.

Next, we have studied the warped spaces of the Klein–Gordon simple separable geometries. In this way we have constructed a more general metric for which it is also possible to solve the Klein–Gordon equation by separation of variables. Although not all of these geometries possesses the principal closed conformal Killing–Yano form, they are equipped with the tower of rank-two Killing tensors and Killing vectors. These symmetry objects allow us to define phase-space variables, which are in involutions and guarantee that geodesic motion is completely integrable. They also define operators which commute with each other and the common set of eigenfunctions has been found by separation of variables.

Finally, we have solved the Einstein equations for the warped metric. We have recovered the spaces which

can be obtained as a limit of vanishing rotations from the Kerr–NUT–(A)dS metric [42]. Beside them we have found spacetimes for which the metric block under the warp factor is the Klein–Gordon simple separable Einstein–Kähler metric. These spacetimes are new solutions in higher dimensions.

### ACKNOWLEDGMENTS

This work was supported by the project of excellence of the Czech Science Foundation No. 14-37086G. I.K. was also supported by the Charles University Grant No. SVV-260211. The authors would like to thank Filip Hejda for valuable discussions on Carter’s paper

### Appendix A: Useful identities

We list here some important identities for functions  $A_\mu^{(i)}$ ,  $U_\mu$  which emerge in the definition of metric (2.1):

$$\sum_\mu \frac{A_\mu^{(i)}}{U_\mu} (-Z_\mu)^{N-1-j} = \delta_{ij} , \quad (A1)$$

$$\sum_i \frac{A_\mu^{(i)}}{U_\nu} (-Z_\nu)^{N-1-i} = \delta_{\mu\nu} ,$$

$$\frac{1}{U_\nu} \sum_{\substack{\mu \\ \mu \neq \nu}} \frac{1}{Z_\mu - Z_\nu} = \sum_{\substack{\mu \\ \mu \neq \nu}} \frac{1}{U_\mu} \frac{1}{Z_\nu - Z_\mu} , \quad (A2)$$

$$A^{(i)} = A_\mu^{(i)} + Z_\mu A_\mu^{(i-1)} , \quad (A3)$$

$$\sum_\mu \frac{A_\mu^{(j)}}{Z_\mu U_\mu} = \frac{A^{(j)}}{A^{(N)}} , \quad (A4)$$

$$U_{\mu,\nu} = \delta_{\mu\nu} \sum_{\substack{\alpha \\ \alpha \neq \mu}} \frac{Z'_\mu}{Z_\mu - Z_\alpha} U_\mu + (1 - \delta_{\mu\nu}) \frac{Z'_\nu}{Z_\nu - Z_\mu} U_\mu , \quad (A5)$$

$$A_{\mu,\nu}^{(i)} = \frac{Z'_\nu}{Z_\nu - Z_\mu} (A_\mu^{(i)} - A_\nu^{(i)}) , \quad (A6)$$

$$\left( \frac{U}{U_\mu} \right)_{,\mu} = 0 , \quad (A7)$$

$$\left( \frac{A^{(i)}}{A^{(N)}} \right)_{,\mu} = - \frac{Z'_\mu}{Z_\mu} \frac{A_\mu^{(i)}}{A^{(N)}} . \quad (A8)$$

We state here an important lemma, which in one direction is a consequence of identities (A1) and in the other direction involves more delicate arguments, cf. [48].

**Lemma A.1** *Functions  $F_\mu = F_\mu(x_\mu)$  satisfy equation*

$$\sum_\mu \frac{F_\mu}{U_\mu} = 0 , \quad (A9)$$

*if and only if they are given by*

$$F_\mu = \sum_{k=0}^{N-2} a_k (-Z_\mu)^k , \quad (A10)$$

*where  $a_k$  are arbitrary constants.*

If the right-hand side of equation (A9) is nontrivial, the solution of is given by a homogeneous solution (A10) plus a nonhomogeneous solution. Two special nonhomogeneous solutions, which follow from identities (A1) and (A4), are

$$\sum_\mu \frac{(-Z_\mu)^{N-1}}{U_\mu} = 1 , \quad \sum_\mu \frac{1}{Z_\mu U_\mu} = \frac{1}{A^{(N)}} . \quad (A11)$$

### Appendix B: Curvature of the Klein–Gordon separable metric

From the first structure equations,

$$\mathbf{d}e^a + \omega^a_b \wedge e^b = 0 , \quad \omega_{ab} + \omega_{ba} = 0 , \quad (B1)$$

and by employing (A5) and (A6), we obtain the connection 1-forms

$$\begin{aligned} \omega_{\mu\nu} &= (1 - \delta_{\mu\nu}) \frac{1}{2} \frac{1}{Z_\nu - Z_\mu} \left( Z'_\nu \sqrt{\frac{X_\nu}{U_\nu}} e^\mu + Z'_\mu \sqrt{\frac{X_\mu}{U_\mu}} e^\nu \right) , \\ \omega_{\hat{\mu}\hat{\nu}} &= (1 - \delta_{\mu\nu}) \frac{1}{2} \frac{1}{Z_\nu - Z_\mu} \left( Z'_\mu \sqrt{\frac{X_\nu}{U_\nu}} e^\mu + Z'_\nu \sqrt{\frac{X_\mu}{U_\mu}} e^\nu \right) , \\ \omega_{\mu\hat{\nu}} &= \delta_{\mu\nu} \left[ - \left( \sqrt{\frac{X_\mu}{U_\mu}} \right)_{,\mu} \hat{e}^\mu + \frac{1}{2} \sum_{\substack{\rho \\ \rho \neq \mu}} \frac{Z'_\mu}{Z_\mu - Z_\rho} \sqrt{\frac{X_\rho}{U_\rho}} \hat{e}^\rho \right] \\ &\quad + (1 - \delta_{\mu\nu}) \frac{1}{2} \frac{Z'_\mu}{Z_\mu - Z_\nu} \left( \sqrt{\frac{X_\nu}{U_\nu}} \hat{e}^\mu - \sqrt{\frac{X_\mu}{U_\mu}} \hat{e}^\nu \right) . \end{aligned} \quad (B2)$$

Straightforward, but very long, calculations of curvature 2-forms  $\Omega^a_b$  from the second structure equations,

$$\Omega^a_b = \mathbf{d}\omega^a_b + \omega^a_c \wedge \omega^c_b \quad (B3)$$

lead to the Riemann tensor  $R = \Omega_{ab} e^a e^b$ ,

$$\begin{aligned}
\mathbf{R} = & \sum_{\substack{\mu,\nu \\ \nu \neq \mu}} \frac{1}{4} V_{\mu\nu} (e^\mu \wedge e^\nu) \otimes (e^\mu \wedge e^\nu) + \sum_{\substack{\mu,\nu \\ \nu \neq \mu}} \frac{1}{2} W_{\mu\nu} (e^\mu \wedge e^\nu) \otimes (\hat{e}^\mu \wedge \hat{e}^\nu) \\
& + \sum_{\substack{\mu,\nu \\ \nu \neq \mu}} \frac{1}{4} \left( V_{\mu\nu} + \frac{1}{4} \frac{S_{\nu,\mu}}{Z'_\mu} \frac{X_\mu}{U_\mu} + \frac{1}{4} \frac{S_{\mu,\nu}}{Z'_\nu} \frac{X_\nu}{U_\nu} \right) (\hat{e}^\mu \wedge \hat{e}^\nu) \otimes (\hat{e}^\mu \wedge \hat{e}^\nu) + \sum_{\substack{\mu,\nu,\kappa \\ \nu \neq \mu \\ \kappa \neq \mu,\nu}} \frac{1}{2} P_{\mu\nu\kappa} \sqrt{\frac{X_\nu}{U_\nu} \frac{X_\kappa}{U_\kappa}} (\hat{e}^\mu \wedge \hat{e}^\nu) \otimes (\hat{e}^\mu \wedge \hat{e}^\kappa) \\
& + \sum_{\substack{\mu,\nu \\ \nu \neq \mu}} \frac{1}{4} \frac{S_{\nu,\mu}}{Z'_\mu} \sqrt{\frac{X_\mu}{U_\mu} \frac{X_\nu}{U_\nu}} (e^\mu \wedge \hat{e}^\mu) \otimes (e^\mu \wedge \hat{e}^\nu) + \sum_{\substack{\mu,\nu \\ \nu \neq \mu}} \frac{1}{2} \left( V_{\mu\nu} + \frac{1}{4} \frac{S_{\mu,\nu}}{Z'_\nu} \frac{X_\nu}{U_\nu} \right) (e^\mu \wedge \hat{e}^\nu) \otimes (e^\mu \wedge \hat{e}^\nu) \\
& + \sum_{\substack{\mu,\nu \\ \nu \neq \mu}} \frac{1}{2} W_{\mu\nu} (e^\mu \wedge \hat{e}^\nu) \otimes (e^\nu \wedge \hat{e}^\mu) + \sum_{\substack{\mu,\nu,\kappa \\ \nu \neq \mu \\ \kappa \neq \mu,\nu}} \frac{1}{2} P_{\mu\nu\kappa} \sqrt{\frac{X_\nu}{U_\nu} \frac{X_\kappa}{U_\kappa}} (e^\mu \wedge \hat{e}^\nu) \otimes (e^\mu \wedge \hat{e}^\kappa) \\
& + \sum_{\substack{\mu,\nu \\ \nu \neq \mu}} W_{\mu\nu} (e^\mu \wedge \hat{e}^\mu) \otimes (e^\nu \wedge \hat{e}^\nu) + \sum_{\mu} \frac{1}{4} \left[ - \left( \sum_{\nu} \frac{X_\nu}{U_\nu} \right)_{,\mu\mu} - \sum_{\substack{\nu \\ \nu \neq \mu}} \frac{1}{2} \frac{S_{\nu,\mu}}{Z'_\mu} \frac{X_\nu}{U_\nu} \right] (e^\mu \wedge \hat{e}^\mu) \otimes (e^\mu \wedge \hat{e}^\mu).
\end{aligned} \tag{B4}$$

Here, for two 2-forms we, define  $\alpha \otimes \beta = \alpha\beta + \beta\alpha$ , and

$$\begin{aligned}
V_{\mu\nu} &= \frac{1}{4} \frac{1}{Z_\nu - Z_\mu} \left[ \frac{Z_\mu}{Z'_\mu} \left( \sum_{\kappa} \frac{Z'_\kappa{}^2}{Z_\kappa} \frac{X_\kappa}{U_\kappa} \right)_{,\mu} - \frac{Z_\nu}{Z'_\nu} \left( \sum_{\kappa} \frac{Z'_\kappa{}^2}{Z_\kappa} \frac{X_\kappa}{U_\kappa} \right)_{,\nu} \right], \\
W_{\mu\nu} &= \frac{1}{4} \frac{1}{Z_\nu - Z_\mu} \left[ Z'_\nu \left( \sum_{\kappa} \frac{X_\kappa}{U_\kappa} \right)_{,\mu} - Z'_\mu \left( \sum_{\kappa} \frac{X_\kappa}{U_\kappa} \right)_{,\nu} \right], \\
P_{\mu\nu\kappa} &= \frac{1}{4} \left[ \frac{Z'_\mu{}^2}{(Z_\mu - Z_\nu)(Z_\mu - Z_\kappa)} + \frac{Z'_\nu{}^2}{(Z_\nu - Z_\mu)(Z_\nu - Z_\kappa)} + \frac{Z'_\kappa{}^2}{(Z_\kappa - Z_\mu)(Z_\kappa - Z_\nu)} \right].
\end{aligned} \tag{B5}$$

- 
- [1] Roy P. Kerr, “Gravitational field of a spinning mass as an example of algebraically special metrics,” *Phys. Rev. Lett.* **11**, 237–238 (1963).
  - [2] Brandon Carter, “Hamilton-Jacobi and Schrödinger separable solutions of Einstein’s equations,” *Commun. Math. Phys.* **10**, 280–310 (1968).
  - [3] Brandon Carter, “Republication of: Black hole equilibrium states,” *General Relativity and Gravitation* **41**, 2873–2938 (2009).
  - [4] F. R. Tangherlini, “Schwarzschild field in  $n$  dimensions and the dimensionality of space problem,” *Nuovo Cimento* **27**, 636 (1963).
  - [5] R. C. Myers and M. J. Perry, “Black holes in higher dimensional space-times,” *Ann. Phys. (N.Y.)* **172**, 304–347 (1986).
  - [6] S. W. Hawking, C. J. Hunter, and M. M. Taylor-Robinson, “Rotation and the AdS/CFT correspondence,” *Phys. Rev. D* **59**, 064005 (1999), arXiv: hep-th/9811056.
  - [7] G. W. Gibbons, H. Lü, Don N. Page, and C. N. Pope, “Rotating black holes in higher dimensions with a cosmological constant,” *Phys. Rev. Lett.* **93**, 171102 (2004), arXiv: hep-th/0409155.
  - [8] G. W. Gibbons, H. Lü, Don N. Page, and C. N. Pope, “The general Kerr-de Sitter metrics in all dimensions,” *J. Geom. Phys.* **53**, 49–73 (2005), arXiv: hep-th/0404008.
  - [9] W. Chen, H. Lü, and C. N. Pope, “General Kerr-NUT-AdS metrics in all dimensions,” *Class. Quantum Grav.* **23**, 5323–5340 (2006), arXiv: hep-th/0604125.
  - [10] W. Chen, H. Lü, and C. N. Pope, “Kerr-de Sitter black holes with NUT charges,” *Nucl. Phys.* **B762**, 38–54 (2007), arXiv: hep-th/0601002.
  - [11] Don N. Page, David Kubizňák, Muraari Vasudevan, and Pavel Krtouš, “Complete integrability of geodesic motion in general higher-dimensional rotating black hole spacetimes,” *Phys. Rev. Lett.* **98**, 061102 (2007), arXiv: hep-th/0611083.
  - [12] Valeri P. Frolov, Pavel Krtouš, and David Kubizňák, “Separability of Hamilton–Jacobi and Klein–Gordon equations in general Kerr-NUT-AdS spacetimes,” *J. High Energy Phys.* **0702**, 005 (2007), arXiv: hep-th/0611245.
  - [13] Artur Serdyukov and Pavel Krtouš, “Complete set of commuting symmetry operators for the Klein-Gordon equation in generalized higher-dimensional Kerr-NUT-(A)dS spacetimes,” *Phys. Rev. D* **77**, 044033 (2008), arXiv: 0711.4623 [hep-th].
  - [14] Marco Cariglia, Pavel Krtouš, and David Kubizňák, “Commuting symmetry operators of the Dirac equation, Killing-Yano and Schouten-Nijenhuis brackets,” *Phys. Rev. D* **84**, 024004 (2011), arXiv: 1102.4501 [hep-th].
  - [15] Marco Cariglia, Pavel Krtouš, and David Kubizňák, “Dirac equation in Kerr-NUT-(A)dS spacetimes: Intrinsic characterization of separability in all dimensions,” *Phys. Rev. D* **84**, 024008 (2011), arXiv: 1104.4123 [hep-th].

- [16] Pavel Krtouš, David Kubizňák, Don N. Page, and Valeri P. Frolov, “Killing–Yano tensors, rank-2 Killing tensors, and conserved quantities in higher dimensions,” *J. High Energy Phys.* **0702**, 004 (2007), arXiv: hep-th/0612029.
- [17] Pavel Krtouš, David Kubizňák, Don N. Page, and Muraari Vasudevan, “Constants of geodesic motion in higher-dimensional black-hole spacetimes,” *Phys. Rev. D* **76**, 084034 (2007), arXiv: 0707.0001 [hep-th].
- [18] Martin Walker and Roger Penrose, “On quadratic first integrals of the geodesic equations for type {22} spacetimes,” *Commun. Math. Phys.* **18**, 265–274 (1970).
- [19] Saul A. Teukolsky, “Rotating black holes - separable wave equations for gravitational and electromagnetic perturbations,” *Phys. Rev. Lett.* **29**, 1114–1118 (1972).
- [20] Saul A. Teukolsky, “Perturbations of a rotating black hole. 1. fundamental equations for gravitational electromagnetic and neutrino field perturbations,” *Astrophys. J.* **185**, 635–647 (1973).
- [21] Brandon Carter, “Killing tensor quantum numbers and conserved currents in curved space,” *Phys. Rev. D* **16**, 3395–3414 (1977).
- [22] M. Demianski and M. Francaviglia, “Type-D space-times with a Killing tensor,” *Journal of Physics A: Mathematical and General* **14**, 173 (1981).
- [23] Valeri Frolov and Dejan Stojković, “Particle and light motion in a space-time of a five-dimensional rotating black hole,” *Phys. Rev. D* **68**, 064011 (2003), arXiv: gr-qc/0301016.
- [24] Valeri Frolov and Dejan Stojković, “Quantum radiation from a 5-dimensional rotating black hole,” *Phys. Rev. D* **67**, 084004 (2003), arXiv: gr-qc/0211055.
- [25] Z. W. Chong, G. W. Gibbons, H. Lu, and C. N. Pope, “Separability and Killing tensors in Kerr-Taub-NUT-de Sitter metrics in higher dimensions,” *Phys. Lett.* **B609**, 124–132 (2005), arXiv: hep-th/0405061.
- [26] Muraari Vasudevan, Kory A. Stevens, and Don N. Page, “Separability of the Hamilton-Jacobi and Klein-Gordon equations in Kerr-de Sitter metrics,” *Class. Quantum Grav.* **22**, 339–352 (2005), arXiv: gr-qc/0405125.
- [27] Hari K. Kunduri and James Lucietti, “Integrability and the Kerr-(A)dS black hole in five dimensions,” *Phys. Rev. D* **71**, 104021 (2005), hep-th/0502124.
- [28] Muraari Vasudevan, Kory A. Stevens, and Don N. Page, “Particle motion and scalar field propagation in Myers-Perry black hole spacetimes in all dimensions,” *Class. Quantum Grav.* **22**, 1469–1482 (2005), arXiv: gr-qc/0407030.
- [29] Muraari Vasudevan and Kory A. Stevens, “Integrability of particle motion and scalar field propagation in Kerr-(Anti) de Sitter black hole spacetimes in all dimensions,” *Phys. Rev. D* **72**, 124008 (2005), arXiv: gr-qc/0507096.
- [30] W. Chen, H. Lü, and C. N. Pope, “Separability in cohomogeneity-2 Kerr-NUT-AdS metrics,” *J. High Energy Phys.* **0604**, 008 (2006), arXiv: hep-th/0602084.
- [31] Paul Davis, “A Killing tensor for higher dimensional Kerr-AdS black holes with NUT charge,” *Class. Quantum Grav.* **23**, 3607–3618 (2006), arXiv: hep-th/0602118.
- [32] Valeri P. Frolov and David Kubizňák, “‘Hidden’ symmetries of higher dimensional rotating black holes,” *Phys. Rev. Lett.* **98**, 011101 (2007), arXiv: gr-qc/0605058.
- [33] David Kubizňák and Valeri P. Frolov, “Hidden symmetry of higher dimensional Kerr-NUT-AdS spacetimes,” *Class. Quantum Grav.* **24**, F1–F6 (2007), arXiv: gr-qc/0610144.
- [34] S. Benenti and M. Francaviglia, “Remarks on certain separability structures and their applications to general relativity,” *Gen. Rel. Grav.* **10**, 79–92 (1979).
- [35] S. Benenti and M. Francaviglia, “The theory of separability of the Hamilton-Jacobi equation and its applications to general relativity,” in *General Relativity and Gravitation, vol. I*, edited by A. Held (Plenum Press, New York, 1980) ch. 13.
- [36] Tsuyoshi Houri, Takeshi Oota, and Yukinori Yasui, “Closed conformal Killing–Yano tensor and Kerr-NUT-de Sitter spacetime uniqueness,” *Phys. Lett.* **B656**, 214–216 (2007), arXiv: 0708.1368 [hep-th].
- [37] Pavel Krtouš, Valeri P. Frolov, and David Kubizňák, “Hidden symmetries of higher dimensional black holes and uniqueness of the Kerr-NUT-(A)dS spacetime,” *Phys. Rev. D* **78**, 064022 (2008), arXiv: 0804.4705 [hep-th].
- [38] Tsuyoshi Houri, Takeshi Oota, and Yukinori Yasui, “Generalized Kerr-NUT-de Sitter metrics in all dimensions,” *Phys. Lett.* **B666**, 391–394 (2008), arXiv: 0805.0838 [hep-th].
- [39] Tsuyoshi Houri, Takeshi Oota, and Yukinori Yasui, “Closed conformal Killing-Yano tensor and uniqueness of generalized Kerr-NUT-de Sitter spacetime,” *Class. Quant. Grav.* **26**, 045015 (2009), arXiv: 0805.3877 [hep-th].
- [40] Yuri Chervonyi and Oleg Lunin, “Killing(-Yano) tensors in string theory,” (2015), arXiv: 1505.06154 [hep-th].
- [41] Tsuyoshi Houri, David Kubizňák, Claude M. Warnick, and Yukinori Yasui, “Local metrics admitting a principal Killing-Yano tensor with torsion,” *Class. Quantum Grav.* **29**, 165001 (2012), arXiv: 1203.0393 [hep-th].
- [42] Pavel Krtouš, David Kubizňák, Valeri P. Frolov, and Ivan Kolář, “Deformed and twisted black holes with NUTs,” (2015), to be published, arXiv: 1511.02536 [hep-th].
- [43] Pavel Krtouš, David Kubizňák, and Ivan Kolář, “Killing–Yano forms and Killing tensors on a warped space,” (2015), to be published, arXiv: 1508.02642 [gr-qc].
- [44] Kazuki Hinoue, Tsuyoshi Houri, Christina Rugina, and Yukinori Yasui, “General wahlquist metrics in all dimensions,” *Phys. Rev. D* **90**, 024037 (2014), arXiv: 1402.6904 [gr-qc].
- [45] Ivan Kolář and Pavel Krtouš, “Weak electromagnetic field admitting integrability in Kerr-NUT-(A)dS spacetimes,” *Phys. Rev. D* **91**, 124045 (2015), arXiv: 1504.00524 [gr-qc].
- [46] David Kubizňák, “On the supersymmetric limit of Kerr-NUT-AdS metrics,” *Phys. Lett.* **B675**, 110–115 (2009), arXiv: 0902.1999 [hep-th].
- [47] Claudio Teitelboim, “Gravitation and hamiltonian structure in two spacetime dimensions,” *Phys. Lett.* **B126**, 41–45 (1983).
- [48] Pavel Krtouš, “Electromagnetic field in higher-dimensional black-hole spacetimes,” *Phys. Rev. D* **76**, 084035 (2007), arXiv: 0707.0002 [hep-th].